arXiv:astro-ph/0007304v1 20 Jul 2000

# *Star Formation Efficiency in the Central 1 kpc Region of Early-Type Spiral Galaxies*

*Akio K. Inoue, Hiroyuki Hirashita*[1] *and Hideyuki Kamaya*

*Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, JAPAN*

`AKI: inoue@kusastro.kyoto-u.ac.jp`

## *ABSTRACT*

It has been reported recently that there are some early-type spiral (Sa–Sab) galaxies having evident star-forming regions which concentrate in their own central 1-kpc. In such central region, is the mechanism of the star formation distinct from that in disks of spiral galaxies? To reveal this, we estimate the star formation efficiency (SFE) in this central 1-kpc star-forming region of some early-type spiral galaxies, taking account of the condition for this 1-kpc region to be self-gravitating. Using two indicators of present star formation rate (H$\alpha$ and infrared luminosity), we estimate the SFE to be a few percents. This is equivalent to the observational SFE in the disks of late-type spiral (Sb–) galaxies. This coincidence may support the universality of the mean SFE of spiral galaxies reported in the recent studies. That is, we find no evidence of distinct mechanism of the star formation in the central 1-kpc region of early-type galaxies. Also, we examine the structure of the central star-forming region, and discuss a method for estimating the mass of star-forming regions.

*Key words :* stars: formation — ISM: structure — galaxies: spiral — galaxies: structure

## 1. INTRODUCTION

When we try to reveal the galaxy evolution along with the theory of star and star-cluster formation, it is very important to examine the star formation activity (SFA) of galaxies. The SFA has been frequently examined by using the star formation rate (SFR), which is the mass turned into stars per unit time. Also, the star formation efficiency (SFE) is an important indicator of the SFA. Here, we define the SFE as being the ratio of the stellar mass in a star-forming region to the gas mass of a parent cloud. This SFE relates more directly than the SFR to the mechanism of turning the interstellar matter into stars. Thus, we focus our attention on the SFE in galaxies.

---

[1] Research Fellow of Japan Society for the Promotion of Science.

In this paper, we are interested in early-type spiral galaxies (Sa–Sab), since they are reported to indicate an interesting property of star formation. The SFA in such galaxies is frequently considered to be lower than that in late-type spiral (Sb–) galaxies (e.g., Kennicutt 1998a). However, there are some early-type spiral galaxies showing evident star-forming activity (e.g., Keel 1983; Keel et al. 1985; Kennicutt et al. 1989; Young et al. 1996; Tomita et al. 1996; Devereux & Hameed 1997; Hameed & Devereux 1999). Interestingly, star-forming regions of such early-type spiral galaxies concentrate in their own central 1-kpc region (Usui et al. 1998). According to Inoue, Hirashita & Kamaya (2000, hereafter IHK00), the SFR of this central 1-kpc star-forming region is about 2 $M_\odot$ yr$^{-1}$, that is, the SFR in early-type spirals is comparable to that in late-disks. In order to obtain a more precise picture of the SFA in those central regions, we should examine their SFE. Thus, the main aim of this paper is to estimate the SFE of the central 1-kpc star-forming region in early-type spiral galaxies.

In order to estimate the SFE of a star-forming region, we must examine both of the stellar mass and the gas mass within this region. We will estimate the stellar mass from the H$\alpha$ luminosity by using the formula of the SFR in Kennicutt (1998a) and an appropriate time-scale for the current star formation. We also estimate it from the *IRAS*-far-infrared (FIR) luminosity via the formula of IHK00 which is reviewed in Appendix A. By the way, when we estimate the mass of molecular gas of the sample galaxies, there is no available complete data set for them, unfortunately. Therefore, we estimate the gas mass of the central star-forming regions, taking account of the condition for these regions to be self-gravitating. This is because if the system were not to be self-gravitating, the cite for the star formation could be disturbed by external gravity, so that the star formation itself would be terminated. Also, we discuss the structure of the central star-forming region, and present a method for estimating the gas mass of star-forming regions.

We compare the estimated SFE with typical observational SFEs in the disk of late-type spiral galaxies. As a result, we will show that the SFE in the central part of early-type spiral galaxies is equivalent to that of the late-type disks. This result may support the universality of the mean SFE of spiral galaxies reported by Young et al. (1996) and Rownd & Young (1999). We might need to expect an universal mechanism for star formation in spiral galaxies.

This paper contains the following sections. First, we examine the structure of the central star-forming region in early-type spiral galaxies in §2. Then, we estimate the SFE of this region in §3. Next, we compare our SFE with other observational ones and discuss the critical surface density for self-gravitational contraction to star formation in §4. Conclusions are summarized in the final section.

## 2. STRUCTURE OF CENTRAL STAR-FORMING REGION

In this paper, we will discuss the physical condition of the central 1-kpc region of the early-type spiral galaxies. Hence, first of all, we review the observational picture of those central regions.

Moreover, in this section, we discuss and examine the structure of the star-forming region within the central 1-kpc. A simple structural model for this region is proposed.

### 2.1. Observational properties

Usui et al. (1998) observed H$\alpha$ and $R$-band photometric images (the field of view was $10' \times 6'$) of nearby 15 early-type spiral (Sa–Sab) galaxies which show relatively high star formation activity (SFA) determined from the ratio of *IRAS*-FIR luminosity to $B$-band luminosity. Some properties of their sample galaxies are tabulated in Table 1. They commented that their observed equivalent widths of the H$\alpha$ emission line agree with those measured by Kennicutt (1983) and Romanishin (1990).

When we consider that H$\alpha$ emission originates from H II regions where massive stars have formed, the half-light radius of H$\alpha$ emission represents a characteristic radius of the spatial distribution of star-forming regions. From Table 1, we find that the half-light radii of H$\alpha$ are about 1 kpc (col. [7]). In addition, we find from column 8 that the radii of H$\alpha$ are much smaller than the optical radii which are one-half of the optical diameter, $D_{25}$, in de Vaucouleurs et al. (1991) (RC3). Thus, we expect that the star-forming regions in such galaxies distribute in their central $\sim$ 1 kpc region and concentrate relatively to their optical stellar light.

Next, let us look at kinematical properties of the central 1-kpc region in spiral galaxies. Recently, Sofue et al. (1999) investigated rotation curves of 50 nearby late-type spiral (Sb–Sc) galaxies, using position-velocity diagrams composed of both observations of CO and H I lines. This method is suited to determine the central rotation curves. According to them, typical spiral galaxies show almost flat rotation curves from their central a few hundreds pc to their outer edge. That is, the rotation of a typical spiral galaxy is differential in its central 1-kpc region. Rubin et al. (1985) examined the rotation curves of early-type spirals via the observation of H$\alpha$ line. Then they concluded that the rotation curves of early-type spirals are similar to those of late-type spirals. Thus, we regard the rotation of the central 1-kpc star-forming region in early-type spiral galaxies as being also differential.

Let us turn to the subject of the gas content in the central region. According to Young & Scoville (1991), who presented a detailed review of radial gas distribution, the molecular gas concentrates in the central several kpc region of spiral galaxies, while the H I gas distributions are rather flat throughout their disk. Moreover, the molecular fraction of hydrogen is almost unity in the central a few kpc of nearby spiral galaxies (Sofue et al. 1995; Honma et al. 1995). Also, a dense molecular disk has been observed in the center of the Galaxy (Sanders et al. 1984). Furthermore, a diffuse coronal gas component has been observed at X-ray in many spiral galaxies (e.g., Read et al. 1997). Thus, we can expect that the gas content of the central 1 kpc region is a mixture of the two components; one of the components is a dense molecular gas and the other is a diffuse hot gas.

### 2.2. Characteristic Length Scale of a Differentially Rotating Medium

To make the following discussion clear, as only a first step, we assume that the components of interstellar medium (ISM) in the star-forming regions are well mixed enough for us to regard these regions as uniform. Hereafter, in this paper, 'uniform medium' or 'homogeneous medium' is sometimes used to denote the well-mixed ISM. Also, we assume that these regions rotate differentially and are established with a global pressure balance of the ISM (Myers 1978). If the mixture of some ISM components is well relaxed to equilibrium via pressure, there is a characteristic sound speed which is defined as being the root of the ratio of the mean pressure to the mean density of the system (e.g., Kamaya & Shchekinov 1998). This definition is possible when the thermal process is very active between the two components (e.g., Kamaya 1999). Moreover, we consider that the radius of this region is larger than the characteristic scale of its self-gravity, the Jeans length, since stars and clouds do form in the regions. Then, we present a formula of the Jeans length for our purpose.

For an infinite and homogeneous gaseous system rotating differentially, Bel & Schatzman (1958) and Genkin & Safronov (1975) have examined the self-gravitational instability of the direction perpendicular to the rotational axis in a cylindrically symmetric geometry. According to them, the dispersion relation of a small perturbation in such a system is expressed by the following,

$$\omega^2 = \frac{4\pi^2 {c_s}^2}{\lambda^2} - 4\pi G\rho + \kappa^2 \,, \tag{1}$$

where $\rho$ and $c_s$ denote the mean density and the characteristic sound speed of this system, respectively, and $\omega$ and $\lambda$ are the frequency and the wavelength of the perturbations, respectively. Moreover, $\kappa$ represents an epicyclic frequency of the rotation of this system and it is defined by

$$\kappa^2 \equiv 4\Omega^2 + R\frac{d\Omega^2}{dR} \,, \tag{2}$$

where $\Omega$ is an angular speed of the system and $R$ is a radius from the rotational axis. If the system has a constant rotational speed, $v_{\rm rot}$ (i.e. a differential rotation), then, $\kappa^2 = 2{v_{\rm rot}}^2/R^2$.

When $\omega^2 < 0$, the system becomes unstable owing to its self-gravity. The minimum wavelength that satisfies this condition is called the Jeans length. Therefore, we obtain the Jeans length, $\lambda_J$,

$$\lambda_J = \frac{2\pi c_s}{\sqrt{4\pi G\rho - \kappa^2}} \,. \tag{3}$$

Now, we consider an uniform medium obeying a global pressure balance. The sound speed, $c_s$, is expressed by ${c_s}^2 \approx p/\rho$ and the global pressure balance is represented by $p/k_{\rm B} \approx a$ (a constant), where $p$ is the pressure of this medium, and $k_{\rm B}$ denotes the Boltzmann constant. Thus, equation (3) is reduced to

$${\lambda_J}^2 = \left(\frac{\pi a k_{\rm B}}{G}\right)\left[\rho\left(\rho - \frac{{v_{\rm rot}}^2}{2\pi G R^2}\right)\right]^{-1} \,. \tag{4}$$

Although we find that $a$ is $10^{3-4}$ K cm$^{-3}$ from Myers (1978), we adopt $a$ as $10^4$ K cm$^{-3}$ in this paper, because the central region seems to have a bit higher pressure than the other regions and hot gas component has several $10^{-3}$ H atoms cm$^{-3}$ as its number density and several $10^6$ K as its temperature (e.g., Read et al. 1997).

### 2.3. Required Mean Density for 1-kpc Scale Contraction

The SFA in the early-type spiral galaxies may be explained by the gravitational contraction of the central 1-kpc region. The radius of a star-forming region, $r_{\rm SF}$, should be more than $\lambda_{\rm J}$ to contract by self-gravity, that is $r_{\rm SF} > \lambda_{\rm J}$. From equation (4), we determine the following condition for its contraction:

$$\rho^2 - A\rho - B > 0\,, \tag{5}$$

where

$$A = \frac{{v_{\rm rot}}^2}{2\pi G R^2} \qquad \text{and} \quad B = \frac{\pi a k_{\rm B}}{G {r_{\rm SF}}^2}\,. \tag{6}$$

Since $\rho > 0$, we obtain the solution of inequality (5):

$$\rho > \frac{A}{2} + \frac{A}{2}\sqrt{1 + \frac{4B}{A^2}}\,. \tag{7}$$

When $r_{\rm SF} \sim 1$ kpc (Usui et al. 1998) and $v_{\rm rot} = 200$ km s$^{-1}$ (a typical value for normal size galaxies in Sofue et al. 1999), we obtain the following relation,

$$\frac{4B}{A^2} \simeq 2.7 \times 10^{-3} \left(\frac{R}{\rm kpc}\right)^4 \ll 1\,, \tag{8}$$

as far as $R \lesssim 3$ kpc, which is satisfied for the typical central star-forming regions in early-type spiral galaxies. Thus, the above condition is reduced to $\rho > A$.

Consequently, we determine a condition of the density for gravitational contraction on 1-kpc scale as a function of the radius, and define a critical density, $\rho_{\rm crit}$,

$$\rho(R) > \frac{{v_{\rm rot}}^2}{2\pi G R^2} \equiv \rho_{\rm crit}\,. \tag{9}$$

Since the mean density of the central star-forming region in early-type spiral galaxies should be more than this critical density and the size of the region is estimated to be about 1 kpc, a minimum mean density required for the contraction is

$$\langle\rho\rangle_{\rm min} = \rho_{\rm crit}(1\ {\rm kpc}) \sim 1.0 \times 10^{-22}\,[{\rm g\,cm^{-3}}]\,, \tag{10}$$

or the corresponding number density is

$$\langle n\rangle_{\rm min} \sim 60\,[{\rm H\,atoms\,cm^{-3}}]\,, \tag{11}$$

when we adopt 200 km s$^{-1}$ as the rotational speed.

### 2.4. Structural Model of Central Star-Forming Region

Someone might think that the mean density estimated in the previous subsection is too large since the main component of the ISM in the central regions is just the hot gas bound in a deep gravitational potential well of these regions. Fortunately, however, as commented in §2.1, there are molecular clouds in the central regions, although the spatial distribution of the molecular clouds are not precisely determined. Hence, we examine if this minimum density of equation (10) or (11) can be reproduced easily by the two components of the ISM. For example, when the number density of the component of molecular gas is $10^3$ H atoms cm$^{-3}$ and that of hot gas is several times $10^{-3}$ H atoms cm$^{-3}$, we reproduce the minimum density if we adopt only 0.06 as the volume filling factor of molecular gas. This small filling factor of 0.06 never contradicts to the observations of molecular clouds in the central regions.

Here, we examine if the two components are well mixed via a dynamical process before the star formation onsets. Fortunately, this is possible, because while the sound crossing time-scale over 1 kpc is about $10^7$ years in hot gas, the gravitational time-scale in the minimum density is at most about $10^7$ years. We want to note here that the overall contraction with the Jean-length is very slow (we should never forget this point), since the growth-rate of the mode with Jeans length is zero theoretically (in a realistic condition, of course, the zero growth-rate breaks due to the variety of disturbances). Thus, the two components can be well mixed when the system begins to contract. The effect of the differential motion can also mix the two components, since the rotation time-scale is also about $10^7$ years. To conclude, our estimate in equations (10) and (11) is very reasonable since it is possible for the two components to be well mixed. The above estimation of time-scales confirms our another assumption that the global pressure balance establishes in this region before the star formation onsets, since the molecular clouds are well mixed into the hot gas.

From the above discussions, we regard that the 1 kpc star-forming regions in the center of early-type spiral galaxies are the regions composed of the mixture of the molecular and coronal gas components. The treatment as a self-gravitational contraction of the mixture in 1-kpc scale becomes suitable when their high mean density with the volume filling factor of the cold gas of $\sim 0.06$ is realized. The minimum density for it can be reached and thus the contraction of the system may occur, when the mixture suffers the radiative cooling. We comment again on the requirement for the system to be self-gravitating. If the self-gravity of the system were to be less important, the cite for the star formation could be disturbed very well by the external gravitational force. As a result, the star formation would be terminated there.

## 3. STAR FORMATION EFFICIENCY

We estimate the star formation efficiency (SFE) in the central 1-kpc region of early-type spiral galaxies showing evident star-forming activity. We define the SFE of a star-forming region as the ratio of the stellar mass newly formed in the region ($M_*$) to its total gas mass ($M_{\rm gas}$). That is,

$$\mathrm{SFE} \equiv \frac{M_*}{M_{\rm gas}}\,. \tag{12}$$

In the following subsections, we estimate both $M_{\rm gas}$ (§3.1) and $M_*$ (§3.2), so that we determine the SFE (§3.3).

### 3.1. Estimation of Gas Mass

We first consider the gas mass of the star-forming region. Now, we want to examine the SFA of the 1-kpc regions. Since the thickness of a typical galactic disk is about 100 pc, we cannot consider that such a star-forming region is a sphere. Therefore, the global structure of the star-forming regions are assumed to be a disk.

In order to estimate the gas mass in such disk-like regions, we must consider their thickness. We mean here the term of the thickness as that of the direction parallel to the rotational axis. When the thickness is denoted by $h_{\rm SF}$, the gas mass of a star-forming region is given by

$$M_{\rm gas} = \int_0^{r_{\rm SF}} 2\pi r\, h_{\rm SF}\rho\, dr\,, \tag{13}$$

where $r$ means the radius of the star-forming region extended to $r_{\rm SF}$ and $\rho$ is its mean density.

Since the star-forming regions are self-gravitating, their thickness is more than the Jeans length. In the direction of the thickness, the term of the epicyclic frequency in equation (3) does not exist. Thus, the thickness of the star-forming regions, $h_{\rm SF}$, should satisfy the following condition:

$$h_{\rm SF} > \sqrt{\frac{\pi {c_s}^2}{G\rho}} = \frac{1}{\rho}\sqrt{\frac{\pi a k_{\rm B}}{G}}\,, \tag{14}$$

where we use again ${c_s}^2 \approx a k_{\rm B}/\rho$, which is obtained under the assumption that the well-mixed ISM obeys a global pressure balance (§2.2).

From inequality (14), we find that the minimum of $h_{\rm SF}\rho$ depends on only $a$, that is,

$$h_{\rm SF}\rho > \sqrt{\frac{\pi a k_{\rm B}}{G}}\,. \tag{15}$$

If we assume that the global pressure ($a k_{\rm B}$) does not vary significantly in the central star-forming regions, we obtain the condition for the gas mass in such star-forming regions as the following,

$$M_{\rm gas} > \pi {r_{\rm SF}}^2 h_{\rm SF}\rho = \pi {r_{\rm SF}}^2\sqrt{\frac{\pi a k_{\rm B}}{G}}\,, \tag{16}$$

where $r_{\rm SF}$ is the radius or the size of a star-forming region. When we adopt $10^4$ K cm$^{-3}$ as $a$, we obtain the following condition;

$$\frac{M_{\rm gas}}{M_\odot} > 1.2 \times 10^8 \left(\frac{r_{\rm SF}}{\rm kpc}\right)^2 . \tag{17}$$

Therefore, we can estimate the minimum gas mass of the central star-forming regions through inequality (17) under the assumption that a global pressure balance is established in this region.

From the observation of Usui et al. (1998), we adopt 1 kpc as the radius of the central star-forming region in early-type spiral galaxies. Therefore, we estimate the characteristic gas mass of such regions to be more than $1.2 \times 10^8 M_\odot$. Since it is reported that the mass of molecular gas within the central 1.5 kpc of the Galaxy is about $5 \times 10^8 M_\odot$ (Sanders et al. 1984), the estimated gas mass may be reasonable. Also, since the central molecular mass for four sample galaxies has been reported, we discuss them individually in §3.4.

## 3.2. Estimation of Stellar Mass

### *3.2.1. From the Hα Luminosity*

The H$\alpha$ luminosity is the most familiar tracer of young massive stars. Thus, we estimate the stellar mass newly born from the H$\alpha$ luminosity. According to Kennicutt (1998a), the formula to estimate the SFR from the H$\alpha$ luminosity, $L_{\rm H\alpha}$, is

$$\frac{\rm SFR}{M_\odot\,{\rm yr}^{-1}} = 3.05 \times 10^{-8} \frac{L_{\rm H\alpha}}{L_\odot} . \tag{18}$$

When a typical time-scale of the star formation, $t_{\rm SF}$, is given and the SFR does not vary significantly during this time-scale, the newly born stellar mass, $M_*$, is estimated to be

$$M_* = {\rm SFR} \times t_{\rm SF} . \tag{19}$$

In this paper, we consider the time-scale, $t_{\rm SF}$, to be a luminosity-weighted average lifetime of the main-sequence OB stars which are representative of the young population, since we are interested in the current star formation rate to obtain the present SFE. Thus, we regard the time-scale of the star formation as a typical main-sequence lifetime of OB stars. IHK00 have estimated the lifetime to be $3.3 \times 10^6$ yr, adopting their mass-luminosity relation, main-sequence lifetime as a function of stellar mass, and Salpeter's IMF. We adopt their value here (i.e., $t_{\rm SF} = 3.3 \times 10^6$ yr). Moreover, we assume the SFR to be nearly constant during this timescale because it is much shorter than the timescale of the galaxy evolution (typically $10^9$ yr). Therefore, the stellar mass expected from the H$\alpha$ luminosity is

$$\frac{M_*}{M_\odot} = 0.10 \frac{L_{\rm H\alpha}}{L_\odot} . \tag{20}$$

The observed luminosities of H$\alpha$ + [N II] line of sample galaxies in Usui et al. (1998) are tabulated in Table 1. Their averaged luminosity is $(3.4 \pm 3) \times 10^7 L_\odot$[2]. To obtain a reasonable luminosity due to only starformation, we correct it for the contamination of [N II] line and the internal extinction (mean $A_{\mathrm{H}\alpha} \sim 0.8$ mag from the private communication with Usui). The corrected H$\alpha$ luminosity is $(4.8 \pm 4) \times 10^7 L_\odot$. Consequently, we estimate the stellar mass to be $(4.8 \pm 4) \times 10^6 M_\odot$.

### *3.2.2. From the Infrared Luminosity*

We can also estimate the stellar mass newly born in galaxies from their infrared (IR) luminosity. There are many formulae to estimate it from IR luminosity as, for example, Kennicutt (1998b). Unfortunately, his formula can be applied to only starburst galaxies because he adopted an assumption that IR luminosity of galaxies is almost equal to their bolometric one. We call this assumption the "starburst approximation." Studies of this kind often adopted this "starburst approximation" (otherwise an *empirical* relation between the luminosities of IR and another wavelength is adopted; for example, Buat & Xu 1996 used the luminosity ratio of IR to ultraviolet). However, this approximation cannot be applied to the galaxies in the sample of Usui et al. (1998). Thus, a new *theoretical* formula to estimate the stellar mass from IR luminosity must be applied to this sample. IHK00 have constructed a new algorithm to do so. Their algorithm is reviewed briefly in Appendix A. The derived formula is

$$\frac{M_*}{M_\odot} = \frac{1.5 \times 10^{-3}(1-\eta)}{0.4 - 0.2f + 0.6\epsilon} \, \frac{L_{\mathrm{FIR}}(40 - 120\mu\mathrm{m})}{L_\odot}\,, \tag{21}$$

where $f$ is a fraction of ionizing photons absorbed by hydrogen atoms in H II regions, $\epsilon$ denotes an efficiency of dust absorption for nonionizing photons, and $\eta$ represents a cirrus fraction. Furthermore, $L_{\mathrm{IR}}^{\mathrm{obs}}(8 - 1000\mu\mathrm{m})$ in equation (A3) is converted to *IRAS*-FIR luminosity, $L_{\mathrm{FIR}}(40 - 120\mu\mathrm{m})$, by the factor of 1.4, assuming that the radiation from dust is the modified black body radiation of 30 K ($I_\nu \propto \nu B_\nu$, where $I_\nu$, $\nu$, and $B_\nu$ are intensity, frequency, and the Plank function, respectively).

The available data set in Usui et al. (1998) has *IRAS*-FIR (40–120 $\mu$m) luminosity of 15 early-type spiral galaxies, and it is tabulated in Table 1. The FIR luminosity of this sample is about $(8.0 \pm 6) \times 10^9 L_\odot$. We estimate a characteristic stellar mass newly formed in the central 1-kpc region of this sample by using this averaged FIR luminosity. Of course, this FIR luminosity originates from the whole galaxy. For the sample of early-type spiral galaxies, however, the SFA in their disks seems to be very low. Thus, after we subtract the "cirrus" component, which originates from dust distributed diffusely in the interstellar space and unrelated to star-forming regions, from the observed FIR luminosity, we assume that the whole of residual FIR luminosity originates in the central 1 kpc star-forming region only.

---

[2] Throughout this paper, the numbers after ± mean the typical dispersions of the sample data

Here, we choose three parameters as the following (see Appendix B): $f = 0.26$, $\epsilon = 0.6$, and $\eta = 0.5$. Since the FIR luminosity of these sample galaxies is $(8.0 \pm 6) \times 10^9 L_\odot$, then, $M_* = (8.5 \pm 6) \times 10^6 M_\odot$. This is consistent with the stellar mass estimated from the H$\alpha$ luminosity. Thus, we have confirmed the applicability of the formula of IHK00.

### 3.3. Estimation of SFE

We now estimate a characteristic SFE of the central 1-kpc star forming region in the sample galaxies of Usui et al. (1998). We have already estimated that the gas mass is heavier than $1.2 \times 10^8 M_\odot$ in §3.1 and that the stellar mass from the H$\alpha$ luminosity is $(4.8 \pm 4) \times 10^6 M_\odot$ in the previous subsection. Therefore, we obtain

$$\mathrm{SFE}_{\mathrm{H}\alpha} < 0.04 \pm 0.03 \,. \tag{22}$$

Moreover, we can estimate the SFE from the FIR luminosity. As seen in §3.2.2, the stellar mass determined by the FIR luminosity is $(8.5 \pm 6) \times 10^6 M_\odot$. Thus, we obtain

$$\mathrm{SFE}_{\mathrm{FIR}} < 0.07 \pm 0.05 \,. \tag{23}$$

Here, we find a good agreement between the both estimations. However, we should note that the way of estimating from the H$\alpha$ luminosity may be more reliable than that from the FIR. This is because the estimation from the FIR luminosity suffers more severe uncertainties, for example, the "cirrus" fraction and the spatial distribution of the FIR luminosity.

Anyway, we conclude that the characteristic SFE expected in the central 1-kpc star forming regions in the early-type spiral galaxies is at most a few percents, which is comparable to SFEs in galactic disks as reviewed in §4.1.

### 3.4. Individual SFE of Four Sample Galaxies

For four of our sample galaxies, the central molecular masses have been reported. In Table 2, the observed molecular masses are tabulated. Thus, we can determine directly their central SFE.

In column (3) of Table 2, we show the radii of the area integrated CO intensity to estimate the molecular mass. These CO radii, $r_{\mathrm{CO}}$, are more extended than those of their star-forming region, $r_{e,\mathrm{H}\alpha}$ in Table 1. Hence, we determine the gas mass of the star forming regions by

$$M_{\mathrm{gas}} = \left(\frac{r_{e,\mathrm{H}\alpha}}{r_{\mathrm{CO}}}\right)^2 M_{\mathrm{H}_2} \,. \tag{24}$$

Consequently, we obtain the following equation from equations (12), (20), and (24);

$$\mathrm{SFE} = 0.10 \left(\frac{r_{\mathrm{CO}}}{r_{e,\mathrm{H}\alpha}}\right)^2 \frac{L_{\mathrm{H}\alpha}/L_\odot}{M_{\mathrm{H}_2}/M_\odot} \,. \tag{25}$$

As shown in column (5) of Table 2, the determined SFEs are about one percent. We find a good agreement between the both SFEs estimated in §3.3 and here. Therefore, our estimation in the previous subsection is quite reasonable.

## 4. DISCUSSIONS

In this section, we discuss some implications derived from our results in the previous sections. Especially, we are interested in whether our SFE is larger or smaller than other SFEs in various conditions of star formation. If our SFE is similar to the mean value, the star formation mechanism, which is typically a mode in molecular clouds, may be universal everywhere, except for starburst, strongly interacting, and so on. Thus, we shall examine the physical reason why our SFE becomes a few percents. For this purpose, we also examine the criterion of the contraction of the parent clouds for star formation.

### 4.1. Comparison with Other Observational SFEs

We compare the SFE estimated in §3.3 with other SFEs determined observationally. Myers et al. (1986) examined 54 molecular clouds associated with H II regions in the Galactic disk. These molecular clouds locates in $-1^\circ \leq b \leq 1^\circ$ and $12^\circ \leq l \leq 60^\circ$. They estimated the molecular mass of each cloud from observations of CO line emission, and also estimated the stellar mass in H II regions accompanying each cloud from H 110$\alpha$ radio (6 cm) emission line. Their result indicates that the averaged SFE is about 0.02. Williams & McKee (1997) also argued that the Galactic averaged SFE is about 0.05. Moreover, the SFEs of two giant H II regions (NGC 604 and NGC 595) in the disk of M33 are reported to be 0.02–0.05 by Wilson & Matthews (1995). Therefore, we find that the SFE expected in the central 1-kpc star-forming region of early-type spiral galaxies is very similar to that in the disk of late-type spiral galaxies.

By the way, we can also define a kind of the SFE of galaxies from the ratio of their luminosity of newborn stars to their molecular mass. When we consider that H$\alpha$ luminosity, $L_{\mathrm{H}\alpha}$, traces the mass of recently formed stars, $L_{\mathrm{H}\alpha}/M_{\mathrm{H}_2}$ is regarded as an indicator of the SFE, where $M_{\mathrm{H}_2}$ is the molecular mass. Young et al. (1996) and Rownd & Young (1999) observed H$\alpha$ and $R$-band luminosities of spiral galaxies in the sample of the Five College Radio Astronomy Observatory (FCRAO) *Extragalactic CO Survey* (Young et al. 1995), and examined morphological variations of the SFE for their sample by using this ratio. Then, they concluded that there is little variation in the mean SFE among the morphologies of the spiral galaxies and it is about 4%. As a result, the coincidence between our theoretical SFE in the center of early-type spiral galaxies and observational ones in the disk of late-type spiral galaxies may support the universality of the mean SFE against the morphological variation of spiral galaxies. This may indicate that the same process such as contraction of molecular clouds by their self-gravity dominates the star formation in all spiral galaxies.

### 4.2. Criterion of Self-Gravitational Contraction

Here, we first discuss inequality (15). The left-hand side of it is the thickness, $h_{\rm SF}$, multiplied by the density, $\rho$, of star-forming regions. This represents a surface density of these regions. Thus, inequality (15) suggests a criterion of the surface density for the self-gravitational contraction on the assumption that the medium in these regions is well mixed and obeys the law of global pressure balance.

We rewrite this criterion of the surface density, defining a critical surface density, $\Sigma_{\rm crit}$;

$$h_{\rm SF}\rho > \sqrt{\frac{\pi a k_{\rm B}}{G}} \equiv \Sigma_{\rm crit}\,. \tag{26}$$

When we adopt $a = 10^4$ K cm$^{-3}$ as the law of the global pressure balance (§2.2), we obtain the following value,

$$\Sigma_{\rm crit} \sim 40\ [M_\odot\,{\rm pc}^{-2}]\,. \tag{27}$$

It is the most important point that the critical surface density depends on only the global pressure represented by $a$. Once $a$ is given, the gas mass of a star-forming region is reduced to a function of its radius through inequality (17).

By the way, the global pressure balance is a balance among various components of the ISM in a steady state. Thus, the ISM evolving dynamically does not always obey this balance. For example, the H II regions, located inside the molecular clouds with active star formation, are not in an equilibrium state but evolve dynamically. Therefore, it is very interesting to examine more precisely how the global pressure balance among the ISM components establishes. Then, we shall compare our critical surface density with an alternative one.

In the 1-kpc star-forming regions, we also consider that the thickness of these regions is more than the Jeans length and the density is also more than the critical density in equation (9). Thus, we find another critical surface density, $\Sigma'_{\rm crit}$;

$$h_{\rm SF}\rho > \sqrt{\frac{\pi\sigma^2\rho_{\rm crit}}{3G}} = \frac{\sigma}{\sqrt{6}\,G}\,\frac{v_{\rm rot}}{R} \equiv \Sigma'_{\rm crit}\,, \tag{28}$$

where we eliminate $\rho_{\rm crit}$, using equation (9), and a characteristic sound speed is rewritten by $(\sigma^2/3)^{1/2}$ in terms of a velocity dispersion, $\sigma$, of the ISM mixture. Therefore, we obtain the following equation,

$$\frac{\Sigma'_{\rm crit}}{M_\odot\,{\rm pc}^{-2}} = \frac{\sigma}{\sqrt{6}\,G}\,\frac{v_{\rm rot}}{R} \simeq 0.6\left(\frac{\sigma}{6.0\ {\rm km\,s^{-1}}}\right)\left(\frac{v_{\rm rot}}{\rm km\,s^{-1}}\right)\left(\frac{\rm kpc}{R}\right)\,, \tag{29}$$

where we have adopted $\sigma \sim 6$ km s$^{-1}$, according to Kennicutt (1989). This is consistent with the critical surface density given in equation (6) of Kennicutt (1989), who derived it from the criterion of instability for a thin isothermal disk of Toomre (1964).

We thus find that the estimated $\Sigma_{\rm crit}$ in equation (27) is nearly equal to $\Sigma'_{\rm crit}$ in equation (29) when we adopt 200 km s$^{-1}$ as $v_{\rm rot}$. Now, we try to find the reason for this coincidence between equations (27) and (29). To determine equation (27), we adopt $p \sim 10^{-12}$ erg cm$^{-3}$, then for $\rho_{\rm crit} \sim 10^{-22}$ g cm$^{-3}$, we find a characteristic sound speed is an order of km s$^{-1}$. By the way, it seems that $\sigma$ is also an order of km s$^{-1}$. This can mean that $\sigma$ is decided by the way how ISM is well mixed and is in global pressure balance. Thus, our critical surface density of equation (27) coincides with equation (29) and with Kennicutt (1989)'s result from the criterion of Toomre (1964), since there exists the law of the global pressure balance in the well-mixed ISM. Anyway, our conclusion is based on the two physical assumptions; (a) there is a global law of the pressure balance among the ISM; (b) the ISM components are well mixed among each other. The first assumption is established via Figure 1 of Myers (1978), and the second one is good as far as the effective crossing time is not longer than the contraction time-scale of the starforing disk as discussed in §2.4.

## 5. CONCLUSIONS

We examine the SFE of the central star-forming region in early-type spiral galaxies, and reach the following conclusions:

[1] We estimate the SFE in the central 1-kpc star-forming region of some early-type spiral galaxies, taking account of the condition for this 1-kpc region to be self-gravitating. The expected SFE is at most a few percents.

[2] The SFE of four sample galaxies (NGC1022, NGC2782, NGC3504, and NGC7625), whose central molecular mass is measured observationally, is estimated to be about one percent.

[3] The estimated SFE is equivalent to the observational ones in the disk of late-type spiral galaxies. This coincidence may support the universality of the mean SFE against the morphological variation of the spiral galaxies.

[4] Assuming a global pressure balance of the ISM, the critical surface density for the self-gravitational contraction depends on the adopted pressure balance. As a result, the mass of star-forming regions is always estimated via only their size once the pressure balance is given.

[5] We suggest that the central 1 kpc star-forming region of early-type spiral galaxies is composed of the mixture of the molecular and coronal gas components. The mixture is possible to contract in 1 kpc scale via their self-gravity if the mean gas density of the mixture is more than a minimum value (e.g., 60 H atoms cm$^{-3}$ in this paper).

[6] The formula of IHK00 is reasonable since SFE estimated from far-infrared luminosity is

nearly equivalent to that determined from H$\alpha$ luminosity.

We would like to thank Prof. M. Saitō for continuous encouragement. One of us (H.H.) acknowledges the Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists.

## A. Newly Formed Stellar Mass Expected from Infrared Luminosity

We present a formula to estimate the stellar mass newly formed in galaxies from their IR luminosity of dust emission. More detailed information can be found in Inoue, Hirashita & Kamaya (2000) (hereafter IHK00) who construct a new algorithm to estimate the SFR from IR luminosity. The result of IHK00 is an extension of a theory of IR radiation from a dusty H II region in Petrosian et al. (1972).

Petrosian et al. (1972) derived a formula of IR luminosity of an H II region, adopting the Case B approximation, which is an assumption of a large optical depth for every Lyman-emission-line photon (Osterbrock 1989), and assuming that all Lyman $\alpha$ photons are absorbed by dust grains in the H II region. Subsequently, IHK00 has developed this result of Petrosian et al. (1972), adopting the Salpeter's IMF, $\psi(m) \propto m^{-2.35}$ for $0.1 M_\odot < m < 100 M_\odot$, a stellar mass-luminosity relation fitted to Table 3.13 of Binney & Merrifield (1998) and a function of the luminosity fraction of ionizing photons from young massive stars fitted to Table 2 in Panagia (1973). They also assume that the luminosity of star-forming regions is dominated by the bolometric luminosities of OB stars on the main-sequence. Moreover, observed IR luminosity of galaxies consists of two components: [1] the component originating from dust nearby star-forming regions and [2] the component originating from dust unrelated to such regions and distributed diffusely in the interstellar space, called the cirrus component (e.g., Helou 1986; Lonsdale-Persson & Helou 1987). Thus, IR luminosity of star-forming regions is the luminosity subtracted this cirrus component from observed IR luminosity. As a result, they obtain the luminosity of star-forming regions,

$$L_{\rm SF} = \frac{1-\eta}{0.4 - 0.2f + 0.6\epsilon}\, L_{\rm IR}^{\rm obs}\,, \tag{A1}$$

where $f$ is a fraction of ionizing photons absorbed by hydrogen atoms in H II regions, $\epsilon$ denotes an efficiency of dust absorption for nonionizing photons and $\eta$ represents a cirrus fraction.

On the other hand, the stellar mass newly formed in star-forming regions expected from their luminosity is expressed by

$$\frac{M_*}{M_\odot} = \frac{\displaystyle\int_{0.1M_\odot}^{100M_\odot} m\psi(m)dm}{\displaystyle\int_{3M_\odot}^{100M_\odot} l(m)\psi(m)dm}\,\frac{L_{\rm SF}}{L_\odot} = 1.1\times 10^{-3}\,\frac{L_{\rm SF}}{L_\odot}\,, \tag{A2}$$

where $m$ denotes a stellar mass in solar unit, $l(m)$ means the adopted mass-luminosity relation as a function of a stellar mass in solar unit, and the mass range of OB stars is adopted 3–$100M_\odot$. From equation (A1) and (A2), IHK00 has obtained the stellar mass expected from observed IR luminosity as the following formula,

$$\frac{M_*}{M_\odot} = \frac{1.1 \times 10^{-3}(1-\eta)}{0.4 - 0.2f + 0.6\epsilon} \frac{L_{\rm IR}^{\rm obs}(8 - 1000\mu{\rm m})}{L_\odot} . \tag{A3}$$

Here, the spectral range of observed IR luminosity is considered to be $8 - 1000\mu$m as the almost whole range of the thermal dust radiation.

Then, IHK00 estimate a typical time-scale of the star formation which is luminosity-weighted average lifetime of the main-sequence OB stars, so that they derive the formula estimating the SFR as the following,

$$\frac{\rm SFR}{M_\odot\,{\rm yr}^{-1}} = \frac{M_*}{t_{\rm SF}} = \frac{3.3 \times 10^{-10}(1-\eta)}{0.4 - 0.2f + 0.6\,\epsilon} \frac{L_{\rm IR}^{\rm obs}(8\text{–}1000\,\mu{\rm m})}{L_\odot} , \tag{A4}$$

where $t_{\rm SF} = 3.3 \times 10^6$ yr as the time-scale of the star formation.

## B. Determination of Three Parameters

Here, we determine values of three parameters in equation (A3). For absorbing fraction by neutral hydrogen, $f$, we adopt 0.26 derived by Petrosian et al. (1972) from an optical depth of dust in Orion nebula. The efficiency of dust absorption for nonionizing photons, $\epsilon$, is estimated to be 0.6 from the averaged 1000–4000Å extinction curve of the Galaxy (Savage & Mathis 1979) and the average visual extinction of Usui's sample ($A_V = 1$ mag from private communication with Usui). Here, we should note that when our formula is applied to the individual molecular clouds, these parameters may depend on the geometry of clouds. Moreover, we choose 0.5 for the cirrus fraction, $\eta$, of Usui's sample, according to Lonsdale-Persson & Helou (1987), who presented a model of the cirrus fraction by using the ratio of IRAS $60\mu$m and $100\mu$m fluxes. Also, we can estimate this fraction by applying a proper radiative transfer model for reproducing multi-wavelength data of galaxies (e.g., Silva et al. 1998, Efstathiou et al. 2000). However, the fraction of the cirrus component remains still uncertain.

## REFERENCES

Bel, N., & Schatzman, E. 1958, RvMP, 30, 1015

Binney, J., & Merrifield, M. 1998, Galactic Astronomy (New Jersey: Princeton University Press) chap.3

Buat, V., & Xu, C. 1996, A&A, 306, 61

de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., Buta, R. J., Patural, G., Fouqeé, P. 1991, Third Reference Catalogue of Bright Galaxies (New York: Springer) (RC3)

Devereux, N. A., & Hameed, S. 1997, AJ, 113, 599

Efstathiou, A., Rowan-Robinson, M., & Siebenmorgen, R. 2000, MNRAS, 313, 734

Garcia-Barreto, J. A., Downes, D., Combes, F., Gerin, M., Carrasco, L., & Cruz-Gonzales, I. 1991, A&A, 252, 19

Genkin, I. L., & Safronov, V. S. 1975, SvA, 19, 189

Hameed, S., & Devereux, N. 1999, AJ, 118, 730

Helou, G. 1986, ApJ, 311, L33

Honma, M., Sofue, Y., & Arimoto, N. 1995, A&A, 304, 1

Inoue, A. K, Hirashita, H., & Kamaya, H. 2000, PASJ, 52, 539 (astro-ph/0003318) (IHK00)

Jogee, S., Kenney, J. D. P., & Smith, B. J. 1999, ApJ, 526, 665

Kamaya, H. 1999, PASJ, 51, 617

Kamaya, H. & Shchekinov, Yu. A., 1998, PASJ, 50, 621

Keel, W. C. 1983, ApJ, 268, 632

Keel, W. C., Kennicutt, R. C. Jr., Hummel, E., & van der Hulst, J. M. 1985, AJ, 90, 708

Kenney, J. D. P., Carlstrom, J. E., & Young, J. S. 1993, ApJ, 418, 687

Kennicutt, R. C. Jr. 1983, ApJ, 272, 54

Kennicutt, R. C. Jr. 1989, ApJ, 344, 685

Kennicutt, R. C. Jr. 1998a, ARA&A, 36, 189

Kennicutt, R. C. Jr. 1998b, ApJ, 498, 541

Kennicutt, R. C. Jr. Edgar, B. K., & Hodge, P. W. 1989, ApJ, 337, 761

Li, J. G., Seaquist, E. R., Wrobel, J. M., Wang, Z. & Sage, L. J. 1993, ApJ, 413, 150

Lonsdale-Persson, C. J., & Helou, G. 1987, ApJ, 314, 513

Myers, P. C. 1978, ApJ, 225, 380

Myers, P. C., Dame, T. M., Thaddeus, P., Cohen, R. S., Silverberg R. F., Dwek, E., & Hauser, M. G. 1986, ApJ, 301, 398

Osterbrock, D. E. 1989, Astrophysics of Gaseous Nebulae & Active Galactic Nuclei (Mill Valley: University Science Books) chap.4

Panagia, N. 1973, AJ, 78, 929

Petrosian, V., Silk, J., & Field, G. B. 1972, ApJ, 177, L69

Read, A. M., Ponman, T. J., & Strickland, D. K. 1997, MNRAS, 286, 626

Romanishin, W. 1990, AJ, 100, 373

Rubin, V. C., Burstein, D., Ford, W. K., & Thonnard, N 1985, ApJ, 289, 81

Rownd, B. K., & Young, J. S. 1999, AJ, 118, 670

Sanders, D. B., Solomon, P. M., & Scoville, N. Z. 1984, ApJ, 276, 182

Silva, L., Granato, G. L., Bressan, A., & Danese, L. 1998, ApJ, 509, 103

Sofue, Y., Honma, M., & Arimoto, N. 1995, A&A, 296, 33

Sofue, Y., Tutui, Y., Honma, M., Tomita, A., Takamiya, T., Hodo, J., & Takeda, Y. 1999, ApJ, 523, 136

Tomita, A., Tomita, Y, & Saitō, M. 1996, PASJ, 48, 285

Toomre, A. 1964, ApJ, 139, 1217

Usui, T., Saitō, M., & Tomita, A. 1998, AJ, 116, 2166

Williams, J. P., & McKee, C. F. 1997, ApJ, 476, 166

Wilson, A. D., & Matthews, B. C. 1995, ApJ, 455, 125

Young, J. S., Allen, L, Kenney, J. D. P., Lesser, A., & Rownd, B. 1996, AJ, 112, 1903

Young, J. S., & Scoville, N. Z. 1991, ARA&A, 29, 581

Young, J. S., Xie, S., Tacconi, L., Knezek, P., Viscuso, P., Tacconi-Garman, L., Scoville, N., Schneider, S., Schloerb, F. P., Lord, S., Lesser, A., Kenney, J., Huang, Y. -L., Devereux, N., Claussen, M., Case, J., Carpenter, J., Berry, M., & Allen, L. 1995, ApJS, 98, 219

---

This preprint was prepared with the AAS LaTeX macros v5.0.

Table 1. Some Properties of Sample Galaxies of Usui et al. (1998)

| Galaxy<br>(1) | $B_T^0$<br>(2) | $\log L_{\rm FIR}$ ($L_\odot$)<br>(3) | $f_{60}/f_{100}$<br>(4) | $\log L_{\rm FIR}/L_B$<br>(5) | $\log L_{{\rm H}\alpha+[{\rm NII}]}$ ($L_\odot$)<br>(6) | $r_{e,{\rm H}\alpha}$ (kpc)<br>(7) | $r_{e,{\rm H}\alpha}/r_{25}$<br>(8) |
|---|---|---|---|---|---|---|---|
| NGC681 | 12.50 | 9.49 | 0.36 | −0.23 | 7.16 | 2.5 | 0.28 |
| NGC1022 | 11.94 | 10.08 | 0.75 | 0.29 | 7.12 | 0.4 | 0.06 |
| NGC2782 | 12.01 | 10.26 | 0.59 | 0.03 | 7.84 | 1.0 | 0.06 |
| NGC2993 | 12.65 | 10.22 | 0.63 | 0.36 | 7.91 | 0.7 | 0.12 |
| NGC3442 | 13.64 | 9.16 | 0.46 | −0.07 | 7.10 | 0.7 | 0.33 |
| NGC3504 | 11.51 | 10.17 | 0.65 | 0.20 | 7.66 | 0.6 | 0.07 |
| NGC3611 | 12.31 | 9.48 | 0.61 | −0.15 | 7.09 | 0.6 | 0.09 |
| NGC3729 | 11.91 | 9.08 | 0.36 | −0.45 | 7.15 | 1.5 | 0.26 |
| NGC4045 | 12.52 | 9.88 | 0.52 | 0.12 | 7.13 | 2.2 | 0.21 |
| NGC4369 | 12.27 | 9.32 | 0.54 | −0.05 | 7.02 | 0.5 | 0.11 |
| NGC4384 | 13.38 | 9.66 | 0.44 | −0.04 | 7.33 | 1.2 | 0.18 |
| NGC5534 | 12.68 | 9.91 | 0.69 | −0.06 | 7.73 | 1.8 | 0.26 |
| NGC5691 | 12.52 | 9.55 | 0.57 | −0.19 | 7.60 | 1.2 | 0.18 |
| NGC5915 | 11.99 | 10.19 | 0.67 | 0.05 | 7.93 | 1.5 | 0.18 |
| NGC7625 | 12.67 | 9.95 | 0.52 | 0.29 | 7.47 | 0.9 | 0.17 |
| average | 12.4±1.1 | $9.9^{+0.2}_{-0.6}$ | 0.6±0.2 | $0.01 \pm 0.2$ | $7.5^{+0.3}_{-0.9}$ | 1.2±0.6 | 0.2±0.2 |

Note. — Col.(1): Galaxy name. Col.(2): $B_T^0$ from the RC3. Col.(3): $\log L_{\rm FIR}$ is calculated using $f_{60}$ and $f_{100}$ from *IRAS* data in solar units. Col.(4): Ratio of $f_{60}$ and $f_{100}$. Col.(5): $\log L_{\rm FIR}/L_B$ is calculated using cols.(2) and (3). Col.(6): Observed H$\alpha$ + [N II] luminosity by Usui et al. (1998). Col.(7): Half-light radius of H$\alpha$ + [N II] emission in kiloparsecs. Col.(8): Ratio of the half-light radius of H$\alpha$ + [N II] emission to the optical radius, $r_{25}$, which is one-half of the optical diameter in the RC3. We assume $H_0 = 75$ km s$^{-1}$ Mpc$^{-1}$. The standard deviation is also shown in the average of each column.

Table 2. Individual Properties of Four Sample Galaxies with CO Observation

| Galaxy (1) | $M_{\rm H_2}$ ($10^9 M_\odot$) (2) | $r_{\rm CO}$ (kpc) (3) | $L_{\rm H\alpha}$ ($10^7 L_\odot$) (4) | SFE (%) (5) | CO Reference (6) |
|---|---|---|---|---|---|
| NGC1022 | 1.5 | 1.65 | 1.8 | 2.0 | Garcia-Barreto et al. (1991) |
| NGC2782 | 2.5 | 1.3 | 9.5 | 0.6 | Jogee et al. (1999) |
| NGC3504 | 1.8 | 0.8 | 6.3 | 0.6 | Kenney et al. (1993) |
| NGC7625 | 2.4 | 1.75 | 4.1 | 0.6 | Li et al. (1993) |

Note. — Col.(1): Galaxy name. Col.(2): Molecular mass estimated from CO intensity via the CO/H$_2$ conversion, $2.8 \times 10^{20}$ H$_2$ cm$^{-2}$/K km s$^{-1}$. Col.(3): Radius of the integrated area to estimate molecular mass. Col.(4): H$\alpha$ luminosity corrected for the contamination of [N II] line and extinction. Col.(5): SFE determined by equation (25).